

latitude  $18^{\circ}50'$  N., longitude  $83^{\circ}20'$  W., reported a barometer reading of 28.79 inches and a wind velocity of 80 miles per hour from the southwest. A later report received by mail from the S.S. *Virginia* which, at the same time, was close to the position of the *President Pierce* gives the following barometer readings: 6 p.m., 29.65 inches; 7 p.m., 29.49 inches; 8 p.m., 28.78 inches; 8:20 to 8:30 p.m. (in calm center, stars visible), 27.44 inches; 9 p.m., 28.64 inches; 10 p.m., 29.24 inches; 11 p.m., 29.40 inches; midnight, 29.70 inches.

This disturbance moved west by north passing inland over the Yucatan Peninsula with center about 40 miles south of Cozumel Island near midnight of the 21st and into the southwestern Gulf of Mexico north of Campeche the evening of the 22d. The center passed inland a short distance south of Tampico, Mexico, the evening of the 24th, attended by winds of hurricane force. The evening report of the 24th received from the S. S. *J. N. Danziger* was remarkable because of the fact that the vessel was at the time in the center of the storm and reported a wind velocity of only 2 miles per hour and a barometer reading of 28.40 inches. As in the case of the

storm of the 15th, great damage was done at and near Tampico, but details are not available.

*Tropical disturbance of September 27–October 4.*—A minor disturbance apparently moved westward between the islands of St. Kitts and St. Martin on the 27th. It was of such small diameter and slight intensity that the center could not be located definitely every 12 hours. However, available data indicate that it moved westward, until the 29th, when it turned to the north and northeast, passing some distance west of Port au Prince, Haiti, the evening of the 29th and centered north of Puerto Plata, Santo Domingo, the morning of October 1. Still of minor intensity, the disturbance then moved northwestward and later north-northwestward until the 4th, after which it could not be located.

*Tropical disturbance of September 28–30.*—This was a very minor disturbance that apparently developed northwest of the Isthmus of Panama and moved northwestward. Its center passed near Cape Gracias the evening of the 28th, then traveled west-northwestward and passed inland south of Belize, British Honduras, the morning of the 30th.

## BIBLIOGRAPHY

C. FITZHUGH TALMAN, in charge of Library

### RECENT ADDITIONS

The following have been selected from among the titles of books recently received as representing those most likely to be useful to Weather Bureau officials in their meteorological work and studies:

#### [American national red cross]

The tornado of May 1, 1933, in Webster and Bienville Parishes, La. Official report of relief operations of the American Red Cross. [Washington, D.C.] 1933. [6 p.] folder. illus. 23 cm.

#### Aristoteles

Nicolaï Cabe . . . In quatvor libros meteorologicorum Aristoteles commentaria, et quæstiones quatvor tomis compræhensa quibus non solum meteorologica . . . Romæ. 1645. 2v. illus. 30 cm. (Typis Hæredum Francisci Corbelletti MDCXLVI.)

#### Cacciatori, Niccolo

De redigendis ad unicam seriem comparibilem meteorologicis ubique factis observationibus conventio proposita et tabulae supputatae ab Equite Nicolao Cacciatore . . . Panormi. 1832. 82 p. engraving, tables. 27½ cm.

#### Calandrelli, [Abbate] Giuseppe

Meteorological correspondence with Johann Jakob Hemmer, S.J., from 1780 to 1787, consisting of 11 autograph letters from the latter, with 11 autograph copies of Calandrelli's answers, with other manuscripts and printed matter . . . 1780–87. 28½ cm. (letter sizes vary).

#### Calandrelli, [Abbate] Giuseppe

Miscellaneous correspondence. 1778–83. 28½ cm. (letter sizes vary).

#### Castellani, Sir Aldo

Climate and acclimatization: Some notes and observations. London, J. Bale, Sons & Danielsson. 1931. viii, 152 p. pl. maps (1 double), diagr. 18½ cm. "References": p. 134–144.

#### Crowe, P. R.

The analysis of rainfall probability. A graphical method and its application to European data. p. 73–91. pl. figs. 25 cm. (Reprinted from the Scottish geographical magazine, v. 39, March 1933.)

#### Florence (Italy). Osserv. Ximeniano dei P. P. Scolopi

Venticinque anni di valori pentadici. Pte. 2. Tensione—umidità—pioggia. 1933. 73 p. pl. 32 cm. (Pubb. Osserv. Ximeniano. no. 145.)

#### Gedda, Mr. le Baron de

Mémoire sur une nouvelle méthode de construire des hygromètres correspondans à points fixes de sécheresse et d'humidité. Ouvrage, qui a remporté le prix proposé par la Société royale des sciences de Copenhague en MDCCCLXXXIII (1783). Copenhague. 1784. 21 p. engr. (fold.) 23½ cm.

#### Little, W. B.

Science and the weather. London. 1933. 155 p. illus. 19 cm.

#### Mazzarón, Anton Giulio

Nozioni elementari di meteorologia ad uso degli aviatori. Milano. Editore Aeronautica. [pref. 1929.] 53 p. illus. (incl. charts). 25 cm. "Bibliografia": p. [54].

#### Meteorologia parallela

Das ist: Courieuse Nachrichten von dem Wetter und den mit silbigem gantz verwunderlich correspondirenden sogenannten Wetter-Gläsern, vom Anfang des Herbsts A. 1710 bis auf den Frühling dieses jetzt lauffenden 1711 Jahrs. Ausz den in Ulm gehabten alltäglichen observationibus zusammen getragen und dem Druck überlassen. 1. Auszfertigung. Frankfurt und Leipzig. [1711.] 32 p. illus., tables. 16½ cm.

#### Mörikofter, W.

Die Abhängigkeit der Sonnenstrahlungsintensität von der Meereshöhe und ihre Konsequenzen für die Organisation der Lichtklimatischen Forschung. 5 p. 23½ cm. (II. Congrès international de la lumière.)

#### Ortiz, Oscar Rivery

Breve estudio del huracán del lro. al 11 de Noviembre de 1932. Habana. 1933. 49 p. charts (1 fold.) 27½ cm. (Boletín hidrográfico. Suplemento no. 1.)

#### Pfaff, C. H.

Über die frühern strengen Winter und über die strengen Winter des achtzehnten Jahrhunderts bis zum Winter von 1776. Die Geschichte der strengen Winter. Erste Abtheilung. Kiel. 1809. xvi, 176 p. 20 cm.

#### Richter, Georg Friedrich

De natalibus fvlminvm tractatvs. physicus accedit appendix. qva. litterae et. observationes qvaedam hvc. pertinentes. Maffei. Lionii Pagliarini. aliorvmqve continentvr. Lipsiae. 1725. 112 p. 16½ cm.

#### Robinson, Gilbert Wooding

Soils: Their origin, constitution, and classification: An introduction to pedology, London, T. Murby & Co.; New York, D. Van Nostrand Co. 1932. xv, 390 p. front. illus. (maps) diagrs. 22 cm. Bibliography at end of each chapter.

## Tandler, T.

De meteoris libris aliisque solertioribus philosophis conscripta, & disputationibus exposita a Tobia Tandlero. Wittebergae, Typis Johann. Gormanni, Anno Christi (1613). 192 p. 15½ cm.

## Venturelli, Lucia

Contatti di masse d'aria calda e d'aria fredda nell'atmosfera in relazione alla situazione barica. Milano. 1933. 24 p. figs. 24 cm. (Excerpt: Bollettino del Comitato nazionale italiano per la geodesia e la geofisica. Seconda serie. Anno 3. N. 1-2, Luglio 1933-XI.)

## Walter, B.

Über den Unterschied in der Blitzgefahr für hart und für weich gedeckte Gebäude . . . Berlin. 1931. 4 p. illus. 30 cm. (Sonderabdruck aus der Elektrotechnischen Zts. 52. Jahrg. 1931, Heft 3, Seite 72.)

## Williamson, Robert M.

Tornado characteristics: The Nashville tornado of March 14, 1933: A brief review of tornadoes in Tennessee. Nashville. 1933. p. 237-248. figs. 23½ cm. (Extr.: Journal Tennessee academy of science. v. 8. no. 3. July, 1933.)

## Winkler, Johann Heinrich

De avertendi fulminis artificio ex doctrina electricitatis disserit atque ad trigam orationvm memoriae Heinrichiana Ridelianae et Seyfertianae sacrarvm qvae in avditorio philosophorum D. XV. Septembris A. R. S. MDCCCLII . . . Lipsiae. 1753. 20 p. front. (engraving). 22½ cm.

## SOLAR OBSERVATIONS

## SOLAR RADIATION MEASUREMENTS DURING SEPTEMBER 1933

By IRVING F. HAND, Assistant in Solar Radiation Investigations

For a description of instruments employed and their exposures, the reader is referred to the January 1932 REVIEW, page 26.

Beginning with this issue, summaries of the total radiation (direct + diffuse) received on a horizontal surface at the Harvard Meteorological Observatory, Blue Hill, Mass. (latitude 42°13' N., longitude 71°07' W.; height above sea level, 195 meters), will be regularly included in table 2. Table 4, giving the values of the red and yellow components of the direct solar radiation at this same station, also will be published regularly, beginning with this number.

Table 1 shows that solar radiation intensities averaged above normal for September at all three Weather Bureau stations.

Table 2 shows an excess in the total solar radiation received on a normal surface at all stations for which normals have been computed except Washington, Lincoln, Chicago, New York, and Pittsburgh. It seems significant that the larger cities show a deficiency in the total radiation received coincident with an increased activity in manufacturing, while the smaller cities show an excess in the radiation received.

In table 3 the turbidity values show a clearing of the sky up to noon and a gradual increase in dustiness during the afternoon. On the 19th the sky gradually cleared up to noon after which cumuli formed.

Polarization measurements made on 4 days at Washington give a mean of 56 percent with a maximum of 64 percent on the 18th. At Madison, measurements made on 9 days give a mean of 68 percent with a maximum of 77 percent on the 27th. These values are slightly above normal for September at both stations.

TABLE 1.—Solar radiation intensities during September 1933

[Gram-calories per minute per square centimeter of normal surface]

## WASHINGTON, D.C.

Date	Sun's zenith distance										
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon
	75th mer. time	Air mass					Local mean solar time				
e.	5.0	4.0	3.0	2.0	1.0 <sup>1</sup>	2.0	3.0	4.0	5.0	e.	
mm	cal	cal	cal	cal	cal	cal	cal	cal	cal	mm	
Sept. 5	12.24		0.90	1.06	1.34					16.20	
Sept. 7	14.10	0.88	0.48	.64	.88	1.24				17.96	
Sept. 8	14.10	.70	.79	.98	1.14	1.39				12.24	
Sept. 12	8.81					1.00				0.14	
Sept. 18	7.87	1.00	1.08	1.22	1.33	1.48				7.29	
Sept. 19	17.37						1.41	1.30			15.11
Sept. 20	5.79	.91	1.01	1.15						7.57	
Sept. 21	7.29	1.02	1.11	1.23	1.37		1.20			7.87	
Sept. 23	10.21					.84	.98			12.24	
Sept. 27	21.26						1.36	1.55	1.32		5.36
Sept. 28	6.50								1.31	1.12	9.47
Sept. 29	6.27					1.00	1.16	1.30	1.25		6.27
Means						.80	1.03	1.18	1.40	1.27	(1.12)
Departures						-.06	±.00	+.01	+.03	+.10	+.10

TABLE 1.—Solar radiation intensities during September 1933—Con.

[Gram-calories per minute per square centimeter of normal surface]

## MADISON, WIS.

Date	Sun's zenith distance										
	8 a.m.	78.7°	75.7°	70.7°	60.0°	0.0°	60.0°	70.7°	75.7°	78.7°	Noon
	75th mer. time	Air mass					P.M.				
e.	5.0	4.0	3.0	2.0	1.0 <sup>1</sup>	2.0	3.0	4.0	5.0	e.	
mm	cal	cal	cal	cal	cal	cal	cal	cal	cal	mm	
Sept. 5	12.24		0.90	1.06	1.34					16.20	
Sept. 7	14.10	0.88	0.48	.64	.88	1.24				17.96	
Sept. 8	14.10	.70	.79	.98	1.14	1.39				12.24	
Sept. 12	8.81					1.00				0.14	
Sept. 18	7.87	1.00	1.08	1.22	1.33	1.48				7.29	
Sept. 19	17.37						1.41	1.30			15.11
Sept. 20	5.79	.91	1.01	1.15						7.57	
Sept. 21	7.29	1.02	1.11	1.23	1.37		1.20			7.87	
Sept. 23	10.21					.84	.98			12.24	
Sept. 27	21.26						1.36	1.55	1.32		5.36
Sept. 28	6.50								1.31	1.12	9.47
Sept. 29	6.27					1.00	1.16	1.30	1.25		6.27
Means						.80	1.03	1.18	1.40	1.27	(1.12)
Departures						-.06	±.00	+.01	+.03	+.10	+.10

## LINCOLN, NEBR.

Sept. 1	15.11	0.52	0.62	0.90	1.08						17.96
Sept. 4	16.79					1.28					16.20
Sept. 5	16.20						1.17				15.65
Sept. 6	15.11						0.95				12.24
Sept. 7	17.96										16.20
Sept. 8	16.20	.71	.82	.97	1.17						14.60
Sept. 16	10.97	.94	1.09	1.20	1.34	1.56					7.29
Sept. 18	15.65	.84	.98	1.12	1.20						19.89
Sept. 19	9.47	.53	.76	1.01	1.27	1.57	1.23	1.02			6.50
Sept. 20	6.76	.91	1.01	1.14	1.29	1.57	1.26	1.09			6.76
Sept. 21	8.18					1.04	1.12				12.24
Sept. 22	7.87						1.36	1.17	1.07		7.57
Sept. 27	6.02	.81	.86	1.09	1.23	1.46	1.27	1.13	1.00		9.47
Sept. 29	9.83					1.01	1.14	1.31	1.47		6.27
Sept. 30	16.20								1.21	1.06	.92
Means						.75	.89	1.03	1.20	1.46	.80
Departures						+.06	±.08	+.09	+.09	+.06	+.05

## BLUE HILL, MASS.

Sept. 2	13.6				1.03	1.11	1.30	1.10	0.78		11.0
Sept. 9	18.0						1.27	1.43	1.24		15.1
Sept. 11	5.6						1.33	1.50	1.20		5.6
Sept. 12	6.3										6.5
Sept. 13	6.8										8.5
Sept. 18	10.6					1.01	1.13	1.47	1.08	-.96	
Sept. 19	8.2						1.02		1.25		7.9
Sept. 22	9.8						1.05				9.1
Sept. 23	8.8						1.11		1.51		7.9
Sept. 27	16.2							1.06			17.4
Sept. 30	7.9						1.14	1.24	1.14		10.6
Means							1.06	1.19	1.44	1.18	(.87)

<sup>1</sup> Extrapolated.